



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

sides $\alpha=0$, $\beta=0$, $\gamma=0$, $\delta=0$, and the *lengths* of the corresponding sides by a , b , c , and d . Let radius of circle be r . Then the equation of the line joining the middle points of the diagonals is $a\alpha - b\beta + c\gamma - d\delta = 0 \dots (1)$ (Cf. Salmon's *Conics*, page 54, Ex. 5).

Putting $\alpha=\beta=\gamma=\delta=r$ we obtain $r(a-b+c-d)=0$, which is satisfied since $a+c=b+d$.

137. Proposed by J. W. YOUNG, Oliver Graduate Scholar in Mathematics, Cornell University, Ithaca, N. Y.

A right cone has its vertex in a horizontal plane, its axis being perpendicular to the plane. A string has one extremity attached to a point on the cone. The other extremity, P , of the string is kept in the plane, and the string is then wound around the cone, without being allowed to slip. Show that the spiral generated by P cuts all straight lines through the vertex at the same angle.

Solution by the PROPOSER.

Let P , P' be two points on the spiral; Q , Q' the corresponding points in the path of the string around the cone; N , N' the points where the perpendiculars from Q , Q' to the plane through the vertex O of the cone, cut the plane.

The right-angled triangles QNO , $Q'N'O$ have the angles QON and $Q'ON'$ equal; hence they are similar.

$$\therefore \frac{QN}{ON} = \frac{Q'N'}{ON'} \dots (1).$$

Again, since the string must not slip, it makes a constant angle with the plane.

$\therefore \triangle QNP$ is similar to $\triangle Q'N'P'$.

$$\therefore \frac{PN}{QN} = \frac{P'N'}{Q'N'} \dots (2).$$

From (1) and (2),


$$\therefore \frac{PN}{ON} = \frac{P'N'}{ON'} \dots (3).$$

But the triangles ONP and $ON'P'$ are right-angled at N and N' (PN , $P'N'$ being the projections to tangents to the circular cone). From (3),

$\therefore \triangle ONP$ is similar to $\triangle ON'P'$.

$\therefore \angle OPN = \angle OP'N'$.

Observing that PN and $P'N'$ are normals to the spiral, the last equation states that the normals make a constant angle with rays through O . Q. E. D.



AVERAGE AND PROBABILITY.

90. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.

During a heavy rain storm a circular pond is formed in a circular field. If a man undertakes to cross the field in the dark, what is the chance that he will walk into the pond? [From Byerly's *Integral Calculus*.]

I. Solution by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Let O be the center of the circular field, and R its radius; C the center of